

AD-A242 869



FTD-ID(RS)T-0731-91

2

FOREIGN TECHNOLOGY DIVISION




DTIC
ELECTE
DEC 2 1991
S C D

EFFECT OF VIBRATION STIMULATION ON THE ULTRASTRUCTURE OF
MECHANORECEPTOR - PACHINIAN CORPUSCLES

by

V.F. Mashanskiy, A. Mirkin, L.N. Vinnichenko



Approved for public release;
Distribution unlimited.

91-16608



91 11 24 009

PARTIALLY EDITED MACHINE TRANSLATION

FTD-ID(RS)T-0731-91

26 September 1991

EFFECT OF VIBRATION STIMULATION ON THE ULTRASTRUCTURE OF
MECHANORECEPTOR - PACHINIAN CORPUSCLES

By: V.F. Mashanskiy, A. Mirkin, L.N. Vinnichenko

English pages: 8

Source: Tsitologiya, Vol. 11, Nr. 12, Publishing House
"Nauka", Leningrad, 1969, pp. 1572-1575

Country of origin: USSR

This document is a machine translation.

Input by: Teara L. Newell

Merged by: Nancy L. Burns

Requester: FTD/TTTR/Lt Billiana Owens

Approved for public release; Distribution unlimited.

Accession For	
NTIS ORIGIN	<input checked="" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A-1	



THIS TRANSLATION IS A RENDITION OF THE ORIGINAL FOREIGN TEXT WITHOUT ANY ANALYTICAL OR EDITORIAL COMMENT. STATEMENTS OR THEORIES ADVOCATED OR IMPLIED ARE THOSE OF THE SOURCE AND DO NOT NECESSARILY REFLECT THE POSITION OR OPINION OF THE FOREIGN TECHNOLOGY DIVISION.

PREPARED BY:

TRANSLATION DIVISION
FOREIGN TECHNOLOGY DIVISION
WPAFB, OHIO.

U. S. BOARD ON GEOGRAPHIC NAMES transliteration SYSTEM

Block	Italic	Transliteration .	Block	Italic	Transliteration
А а	<i>А а</i>	A, a	Р р	<i>Р р</i>	R, r
Б б	<i>Б б</i>	B, b	С с	<i>С с</i>	S, s
В в	<i>В в</i>	V, v	Т т	<i>Т т</i>	T, t
Г г	<i>Г г</i>	G, g	У у	<i>У у</i>	U, u
Д д	<i>Д д</i>	D, d	Ф ф	<i>Ф ф</i>	F, f
Е е	<i>Е е</i>	Ye, ye; E, e*	Х х	<i>Х х</i>	Kh, kh
Ж ж	<i>Ж ж</i>	Zh, zh	Ц ц	<i>Ц ц</i>	Ts, ts
З з	<i>З з</i>	Z, z	Ч ч	<i>Ч ч</i>	Ch, ch
И и	<i>И и</i>	I, i	Ш ш	<i>Ш ш</i>	Sh, sh
Й й	<i>Й й</i>	Y, y	Щ щ	<i>Щ щ</i>	Shch, shch
К к	<i>К к</i>	K, k	Ъ ъ	<i>Ъ ъ</i>	"
Л л	<i>Л л</i>	L, l	Ы ы	<i>Ы ы</i>	Y, y
М м	<i>М м</i>	M, m	Ь ь	<i>Ь ь</i>	'
Н н	<i>Н н</i>	N, n	Э э	<i>Э э</i>	E, e
О о	<i>О о</i>	O, o	Ю ю	<i>Ю ю</i>	Yu, yu
П п	<i>П п</i>	P, p	Я я	<i>Я я</i>	Ya, ya

*ye initially, after vowels, and after Ъ, Ь; e elsewhere.
When written as ѐ in Russian, transliterate as yě or ě.

RUSSIAN AND ENGLISH TRIGONOMETRIC FUNCTIONS

Russian	English	Russian	English	Russian	English
sin	sin	sh	sinh	arc sh	\sinh^{-1}
cos	cos	ch	cosh	arc ch	\cosh^{-1}
tg	tan	th	tanh	arc th	\tanh^{-1}
ctg	cot	cth	coth	arc cth	\coth^{-1}
sec	sec	sch	sech	arc sch	sech^{-1}
cosec	csc	csch	csch	arc csch	csch^{-1}

Russian English

rot curl
lg log

GRAPHICS DISCLAIMER

All figures, graphics, tables, equations, etc.
merged into this translation were extracted
from the best quality copy available.

Page 1572.

EFFECT OF VIBRATION STIMULATION ON THE ULTRASTRUCTURE OF
MECHANORECEPTOR - PACHINIAN CORPUSCLES.

V. F. Mashanskiy, A. Mirkin and L. N. Vinnichenko.

Laboratory of microscopy of the institute of cytology of the AS USSR
and the institute of physiology of the AS USSR, Leningrad.

Ultrastructural changes in the nervous ending of mechanoreceptors of the Pachinian muscles following adequate vibration treatment have been investigated. A formation of vesicles similar to synaptic ones bordering mitochondria was seen. The significance of these vesicles in the process of adaptation is suggested. Changes in the ultrastructure of mitochondria were traced at early stages of irritation, these changes may be involved in the synthesis of macroergic substances.

Up to now investigations of ultrastructure of mechanoreceptors - Pachinian corpuscles - were conducted on unexcited receptors (Pease and Quilliam, 1957; Mirkin, etc., 1967), while changes, which occur during adequate stimulation, remained without attention. Earlier it was established that the most adequate form of stimulation of the Pachinian corpuscles is vibration (Mirkin, etc., 1967). It was shown on the basis of the observations of cyclic variations in the threshold excitation that the Pachinian corpuscles possess resonance properties (Mirkin, 1966).

Material and procedure.

Experiments were conducted at 25°C on Pachinian corpuscles,

obtained from mesentery of cat. One group of receptors (control) was maintained for 15 min. in a Krebs-Henseleit solution, after which it was fixed, another - after 10-minute rest underwent vibration stimulation with a frequency of 120 Hz and the force of 150 dynes. The time of stimulation for different receptors was 5, 10, 20 or 40 s. After action the material was immediately fixed with 2.5% solution of glutaraldehyde on a phosphate buffer (pH 7.2) at 0° for 1 hour with the subsequent additional fixation by 2% solution of tetroxide tetroxide of osmium on the phosphate buffer for 40 min. After dehydration in alcohols the Pachinian corpuscles poured into the alardite.

Results.

Fig. 1 (see inclusively VII) shows transverse section of nerve end of control receptor. Nerve end has a section of elliptic form, its longitudinal axis is oriented in the direction of "fissure" in the internal bulb. The plasmatic membrane of nerve end forms the folds, which convert into the large apophyses near the "fissure". Cytoplasm contains the numerous proto-neurofibrils, which have the form of dark granules with size of 50 Å or short cylinders in the transverse section. This testifies about the orientation of proto-neurofibrils longitudinal with respect to the nerve end. Mitochondria are located predominantly along the periphery of nerve end near the plasmatic membrane, but their greatest accumulation is observed near the "fissure". Are there located scarce canals and cisterns of smooth

endoplasmic network and single bright vesicles with diameter 250-300 Å.

Fig. 2 (see inclusively VII) presents section of nerve end after vibration stimulation for 5 s. Is observed bloating of separate mitochondria; in such mitochondria the matrix becomes electron-transmissive.

Page 1573.

Fig. 3 and 4 (see inclusively VII) show section of nerve end after vibrations for 10 s. It is evident that the thickness of proto-neurofibrils increases to 80 Å. Impression is created, that the electron density of proto-neurofibrils somewhat grows. Virtually all mitochondria are surrounded by numerous vesicles with a diameter of 250-280 Å, the distances between which are commensurate with their quantity. They are located 100-300 Å from the membranes, which limit mitochondria. Separate vesicles are detected near the plasmatic membrane, but they never are observed in the cytoplasm of cell-satellites of internal bulb or external capsule. Vesicles around the mitochondria are observed and after 20-per-second vibrations (Fig. 5; see inclusively VIII), but, as a rule, were absent after 40-per-second stimulations (Fig. 6). It is necessary to note that in the rare cases the vesicles around the mitochondria were visible after 5- and 40-per-second vibration stimulations, that, possibly, points to the high lability of the described vesicular structures.

As is known, processes of onset and propagation of excitation are accompanied by formation and transportation of cholinergic mediators, included in number of cases into synaptic vesicles (Whittaker et al, 1964; Gray and Guillery, 1966). Similar vesicles were described in dendrites (Blacksted and Kjaerjeim, by 1961; Elfvin, 1963), the gustatory kidneys (Trujillo-Cenoz, 1957), the hair cells of organ of Corti (Vinnikov, 1966), to the axoplasm of the Pachinian corpuscles (Pease and Quilliam, 1957), especially in the region of the "fissure" (Mirkin, etc., 1967). However, these vesicles were detected irregularly, also, in a small quantity.

It was of interest to explain, are discovered vesicles by carriers of cholinergic mediators. In view of the great difficulties of the direct detection of mediators in the Pachinian corpuscles we attempted to detect in the nerve ends the enzymes, which esterify the mediators indicated. For this purpose histochemical investigation with the aid of a light microscope was carried out. Was discovered, just as in the works of Portugalov (1955) and Ulumbekov (1964), the appearance of a reaction product within the limits of nerve end. On the basis of this (in accordance with the representations about nature of the receptor surface as about the peculiar formation of the "extended synapse" - (Shabadash, 1968) it is possible to assume that the vesicles discovered by us are similar to the synaptic.

Previously was discussed possibility of onset of vesicles in synaptic terminations from internal membranes of mitochondria

(Dyachkova, etc., 1962). The arrangement of vesicles in immediate proximity with the mitochondria can be the indirect confirmation of a connection of these two structures. Possibly, the described vesicles appear due to budding, similar to pinocytosis.

Obtained by us data show that shaping of vesicles occurs extremely rapidly. Vesicles around the mitochondria were observed after only 5s in some experiments. About the same is testified that never could not meet the vesicles during their budding; they were detected always more narrowly with those formed. On the other hand, the onset of vesicles continues relatively not long. The vesicles usually were not encountered after 40 s. of vibration action. Probably, this phenomenon to some degree explains the mechanism of the adaptation, which appears as a result of depletion of the ability of mitochondria to form vesicles. For a while is required, in order to supplement the supply of mediators, and possibly, also materials for the formation of the membranes around the vesicles. Electrophysiological investigations showed that after 40-per-second stimulations the pulse activity becomes noticeably less frequent (Fig. 7). After certain leisure the activity is restored.

Page 1574.

In early stages of vibration stimulation (5 s.) is discovered bloating of mitochondria. Earlier it was shown (Mashanskiy and Rogozkin, 1965), that an increase of cellular activity is accompanied by bloating of mitochondria, during which is facilitated the

metabolism of the macroergic connections, in particular necessary for the creation of electric potential in the nerve end.

Resume.

Is studied change in ultrastructure of nerve end of Pachinian corpuscles after adequate vibration stimulation. The appearance of vesicles of the type of synaptic, that lie is discovered by border around the mitochondria. In connection with the rapid onset and disappearance of the vesicles indicated is presented the hypothesis about their value in the process of adaptation. In the early stages of excitation are discovered changes in the ultrastructure of the mitochondria, which can be connected with the process of the formation of macroergic compounds.




Fig. 7. Recording of response of receptor during vibration stimulation with frequency 120 hertz. a) after 10 s, stimulation, b) 20, c) 30, d) 40 s, below - time mark.

Page 1575.

REFERENCES.

REFERENCES.

- Винников Я. А. 1966. Структурные и цитохимические основы механизма функции рецепторов органов чувств. В сб.: Нервная клетка, Л.: 7—101. — Дьячкова Л. Н., Давыдова Т. В. и Якобсон И. К. 1962. Участие митохондрий в образовании симпатических пузырьков. ДАН СССР, 147, 6: 1467—1469. — Машанский В. Ф. и Рогожкин В. А. 1965. Изменения ультраструктуры организации митохондрий при мышечной деятельности и гипотермии. В сб.: Электронная и флуоресцентная микроскопия клетки, М.—Л.: 61—67. — Миркин А. С. 1966. О некоторых физических свойствах структуры одиночных рецепторов — телец Пачини. ДАН СССР, 141: 258—261. — Миркин А. С., Тучков Б. С. и Машанский В. Ф. 1967. Некоторые особенности ультраструктуры телец Пачини. Цитология, 9, 6: 637. — Португалов В. В. 1955. Очерки гистологии нервных окончаний. — Улумбеков Э. Г. 1964. О некоторых дегидрогеназах микрокапсулированных нервных окончаний. Арх. анат. гистол. эмбриол., 47, 12: 89—92. — Шабаш А. Л. 1968. Особый коаксиальный нервный аппарат мякотного нервного волокна и роль в проведении импульса. В сб.: Матер. симп. «Некоторые физико-химические акты возбуждения и проведения», М.: 28—29. — Blacksted T. W. and Merzheim A. 1961. Special axodendritic synapses in the hippocampal cortex: electron and light microscopic studies on the layer of mossy fibers. J. comp. neurol., 133—160. — Elfvin L. G. 1963. The ultrastructure of the superior cervical sympathetic ganglion of the cat. II. The structure of the preganglionic end fibers and the axons as studied by serial sections. J. ultrastr. res., 8: 441—476. — Gray E. G. 1966. Synaptic morphology in the normal and degenerating visual system. Intern. rev. cytol., 49: 111—182. — Pease D. C. and Quilley T. A. 1957. Electron microscopy of the Pachinian corpuscle. J. biophys. biochem. opt., 2: 331—342. — Trujillo-Ceno O. 1957. Electron microscope study of rabbit gustatory bud. Zs. Zellforsch., 46: 272—280. — Whittaker V. P., Chaelson I. A. and Kirkland R. J. A. 1964. The separation of synaptic vesicles from nerve-ending particles («Synaptosomes»). Biochem. j., 90: 293—303.

Submitted 12 Apr. 1969.

DISTRIBUTION LIST

DISTRIBUTION DIRECT TO RECIPIENT

ORGANIZATION

MICROFICHE

B085 DIA/RTS-2FI
C509 BALL0C509 BALLISTIC RES LAB
C510 R&T LABS/AVEADCOM
C513 ARRADCOM
C535 AVRADCOM/TSARCOM
C539 TRASANA
Q592 FSTC
Q619 MSIC REDSTONE
Q008 NTIC
Q043 AFMIC-IS
E051 HQ USAF/INET
E404 AEDC/DOF
E408 AFWL
E410 ASDTC/IN
E411 ASD/FTD/TTIA
E429 SD/IND
P005 DOE/ISA/DDI
P050 CIA/OCR/ADD/SD
1051 AFIT/LDE
CCV
PO90 NSA/CDB
2206 FSL

1
1
1
1
1
1
4
1
1
1
1
1
1
1
1
1
2
1
1
1
1

Microfiche Nbr: FTD91C000636
FTD-ID(RS)T-0731-91